

American *Journal of* Pharmacy

AND THE SCIENCES SUPPORTING PUBLIC HEALTH



DR. IVOR GRIFFITH RECEIVING
THE 1960 REMINGTON MEDAL

Since 1825

December 1960

Prepare for a Career in Bacteriology, Biology, Chemistry or Pharmacy

Young men and young women who are interested in productive, satisfying and successful futures in any of these four fields may prepare for ever increasing opportunities through courses of study leading to the B.Sc. degree at this institution, oldest of its kind in the Americas. Selected graduate studies lead to M.Sc. and Ph.D. degrees. Residence Hall for women students now available. Write for 80-page catalog. Terms commence each September.



Philadelphia College of Pharmacy and Science

43d Street, Kingessing and Woodland Avenues

Philadelphia 4, Pa.

Founded 1821

American Journal of Pharmacy

Published monthly by the Philadelphia College of Pharmacy and Science
43d Street, Kingessing and Woodland Avenues, Philadelphia 4, Pa.

Annual Subscription \$4.00
Single Numbers, 40 Cents

Foreign Postage, 25 Cents Extra
Back Numbers, 50 Cents

Second Class Postage Paid at Philadelphia, Pa.



THE CHILLING TALE OF THE PHARMACIST WHOSE ASSETS GOT FROZEN

When the first bargain buy arrived, he made mental calculations and began thinking about Bermuda.

Fifty dozen at \$1.40 each . . . I sell for \$2.75 . . . take out 30 percent for overhead . . . enough left for plane fare . . .

By the time the tenth bargain buy arrived, he was figuring on spending at least two weeks on the French Riviera.

Let's see . . . this year my "buys" will cost me \$25,000, but they'll bring at least \$50,000 at retail. My net will be at least \$10,000. We'll go in January!

But, to his sorrow, most of the bargain merchandise stayed on his shelves, and he spent January wishing he could have a clearance sale like other stores.

Moral:

There's no guaranteed profit in stock. Only sales make money. The Lilly policy of distribution protects the pharmacist from oversupplies of slow-moving items. He buys just what he needs from a Lilly service wholesaler.

If your assets are acquiring a "chill," warm up your operation by routing your orders through one of the 300 Lilly service wholesalers who serve the nation.




ELI LILLY AND COMPANY • INDIANAPOLIS 6, INDIANA, U.S.A.

080957



put
YOURSELF
 in the
 Upjohn
 picture...



The scientists who search for new drugs . . . the chemists who synthesize new compounds . . . the pharmacists who design and prepare the dosage forms . . . the clinicians who show that new agents work where none worked before. The results of all their efforts—vital new drugs for better health—depend for their presentation to the medical profession on a select handful of men. To physicians these professional representatives *are* The Upjohn Company. Physicians turn naturally to them to keep up with advances in Upjohn therapy.

The responsibility is great. Such a man must be well versed in the sciences, with a thorough knowledge of pharmacology and therapeutics; and he must be able to meet members of the medical profession on equal terms. It is quite possible that your pharmaceutical training has equipped *you* for a sales career with Upjohn. If it has, then your future is unlimited. If you feel you are qualified to be trained for one of the most challenging careers the pharmaceutical field has to offer, write for more information to Don Meredith, The Upjohn Company, Kalamazoo, Michigan.

Upjohn

The Upjohn Company
 Kalamazoo, Michigan

AMERICAN
JOURNAL OF PHARMACY
AND THE SCIENCES SUPPORTING PUBLIC HEALTH
Since 1825

LINWOOD F. TICE, Ph. G., M. Sc., D. Sc., Editor
Kenneth Avis, M. Sc., D. Sc., Editorial Assistant
Margaret Mary Culin, Editorial Assistant
John E. Kramer, B. Sc., Business Manager

COMMITTEE ON PUBLICATION

Louis Gershenfeld, P. D., Ph. M., D. Sc., Chairman
Mitchell Bernstein, P. D., M. D., F. A. C. P.
E. Fullerton Cook, P. D., Ph. M., D. Sc.
Marin S. Dunn, A. M., Ph. D.
Joseph W. E. Harrison, P. D., Sc. D.
Ivor Griffith, P. D., Ph. M., D. Sc., F. R. S. A., ex officio

Vol. 132

DECEMBER 1960

No. 12

CONTENTS

Our Cover

- The 1960 Remington Medalist and His Co-Workers.
By L. Gershenfeld 428

Editorial

- The Expansion of Drug Certification by the FDA 431

Article

- Stability Problems With Some Vitamins in Pharmaceu-
ticals. By T. J. Macek 433

- Book Review** 456

- Index to Volume 132** 457

O U R C O V E R

THE 1960 REMINGTON MEDALIST AND HIS CO-WORKERS *

By Louis Gershenfeld, D.Sc.**

TOASTMASTER Doctor Sica, Remington Medalist for 1960, Honored Guests, Members and Friends of the New York Branch of the American Pharmaceutical Association:

It has become my agreeable lot to be the voice of the members of the Faculty of the Philadelphia College of Pharmacy and Science and of other co-workers in the health sciences, who have been associated with the 1960 Medalist during the past several decades. In this capacity for the moment, I cannot help but think of the co-ed who accidentally dropped her expensive camera in the water of a swimming pool adjacent to the campus. She called to one of the professors nearby and asked him to kindly retrieve it. "Why was I chosen?" he asked. "Oh, you don't recognize me, professor" was the reply. "I am in one of your classes and, frankly, I know of no one who can go down deeper, stay down longer, and come up drier than you." Speaking for myself, the time allotted will not permit me to go down too deep, nor stay too long, and I will try not to come up too dry.

There are many descriptive words which can be used describing the Medalist's varied activities during his long, colorful, and productive career. Any one of these alone is enough to make up the life work of most individuals, but to find them all so capably expressed by one individual is extraordinary indeed. A well-known scientist, an inspiring teacher, educator, author, and editor; a capable leader and master executive; a warm friend; and one always cooperating with and solicitous of his co-workers. He is unique in his unselfish service and devotion. The extreme simplicity of his nature, his energy and courage even under adversity, and his boundless en-

* Delivered at a special meeting of the New York Branch, American Pharmaceutical Association, and Testimonial Dinner tendered Doctor Ivor Griffith, recipient of the 1960 Remington Honor Medal, at Hotel Roosevelt, New York City, December 7, 1960.

** Chairman of the Faculty Council and Director of the Department of Bacteriology, Philadelphia College of Pharmacy and Science.

thusiasm and zest for living have earned him universal respect and admiration. He has the capacity to raise men's hopes, strengthen their wills, and bring out the best that is within them. Perhaps the reason for the Medalist's influence on his co-workers can be summed up in the words of Emerson: "This is the key to the power of the greatest men—their spirit diffuses itself."

The pharmaceutical and allied scientific community in this great land of ours grew up during the period of the Medalist's active life. For more than fifty years, he has been associated with the Philadelphia College of Pharmacy and Science—as student, alumnus, President of its Alumni Association, instructor, professor, head of the Department of Pharmacy, dean, emeritus professor, editor of its *American Journal of Pharmacy and the Sciences Supporting Public Health*, the first recipient of the Alumni Award, and for the last twenty years as our Comrade-in-Chief, President of the College. He is the only individual in the one hundred forty years' existence of our College to occupy all of these stations. In associated fields in the health sciences, the 1960 Remington Medalist served in our city as Director of the Pharmaceutical and Clinical Laboratories at Stetson Hospital, as Research Director of the John B. Stetson Co. and also the McNeil Laboratories, as Professor and now Honorary Professor of Organic Chemistry at the Wagner Free Institute of Science, as a former member of the Pennsylvania State Department of Health and just recently appointed for the nineteenth year as Chairman of the State Advisory Committee on Laboratory Procedures, and also as a former Chairman of the Food and Drug Committee of the Philadelphia Department of Public Health. In his duties associated with the John B. Stetson Co., and later as Research Director of the Frank H. Lee Co., at Danbury, Conn., he was successful in replacing the toxic mercurials used in the industry by other more efficient chemicals, thus contributing to the health of the workers in the hat industry. Time is not available to do more than mention the services and his accomplishments as President of the Welsh Society and the St. David's Society as well as his activities in many communal, fraternal, and scientific organizations or to detail the many honors which have come to him. This, however, is worthy of note. After the award of the Remington Honor Medal later this evening, Dr. Griffith will be the only one to have received both the Procter and the Remington Medals, the highest awards in American and probably World Pharmacy.

His intimate friends and co-workers know him as IVOR, the

- I standing for incomparable, informative, and inspiring
- V standing for versatile, vigorous, and vitality
- O standing for organizer, originality, and optimist
- R standing for realist, reliable, and revered.

It is these intimate friends and co-workers who are delighted to be here this evening to extend congratulations and to which we add our tribute. With deep affection and admiration, we join in the universal prayer that our Medalist continue his dedicated life of fruitful service for many, many years to come.

May I conclude by saying we may well note that Doctor Griffith's spirit of self-sacrifice for the good of others is worthy of meditation in all our hearts, and his devotion to his friends and co-workers may well serve as an example for others to emulate?

EDITORIAL

THE EXPANSION OF DRUG CERTIFICATION BY THE FDA

ONE of the several recommendations made by the Committee appointed to study the Food and Drug Administration was that the present program involving the certification of certain antibiotics be expanded to include certification of all antimicrobial substances. Such a recommendation received the endorsement of Secretary Flemming but we sincerely hope that his successor, Mr. Ribicoff, will study this matter more carefully and come to a different conclusion.

The initial reason for the certification of antibiotics was a valid one but the validity of this requirement has long since ceased to exist and it remains with us largely we suspect because the Food and Drug Administration is loath to relinquish what has been a very important source of income for their laboratories outside of their appropriation from Congress.

In the early days, antibiotics were usually quite impure; they varied greatly from batch to batch and it was difficult, if not impossible, to write adequate standards whereby their purity and strength might readily be determined. Under these circumstances, it became necessary for the Food and Drug Administration to examine very carefully each batch of antibiotics produced as the only means available to give the public that protection for which it had the responsibility. Antibiotics today, however, have long since come of age. We look upon these substances now in the same way exactly as we consider hundreds of other potent, life-saving substances. Modern methods of protection and purification have resulted in the preparation of antibiotics which are almost essentially pure substances and, indeed, in most instances, they are recrystallized and possess a degree of purity of about the same order as any other fine chemical. Those responsible for the official compendia; namely, the *United States Pharmacopeia* and the *National Formulary*, prepare and publish very exact standards for these substances, as well as clearly defined procedures which make it possible to determine conformity to these standards. The point which we are making is that there is no real difference between an antibiotic and, shall we say, an alkaloid or glycoside. Each is a very potent and useful substance and one is no more life-saving than another, the determining factor in each case being the disease from which the patient suffers and the medicament

which is needed. Surely, digitalis or digitoxin is just as essential to the patient with congestive heart failure as is an antibiotic to a person suffering with some infection. It is our firm opinion that no clear case can be made for placing antibiotics in a special class and that they should not be so considered.

There are some who might argue that all drugs, regardless of category, should be subjected to precertification by the Food and Drug Administration before they can be sold. By no stretch of the imagination could this be done and, even if it were done, the increased cost—which, of course, would need to be passed to the ultimate consumer—would be staggering. In fiscal 1959, it is claimed, the certification of six antibiotics cost the drug industry 1.2 million dollars. One can only guess what the astronomical figure would be if all drugs were precertified. It has, furthermore, not been demonstrated that certification even of these is necessary. In fact, we suspect that the records would show the reverse to be true.

There are changes needed in the operations of the Food and Drug Administration and these have been well documented for a long time. There needs to be a better inspection of production and control facilities among drug manufacturers, and there needs to be greater effort expended in determining whether drugs shipped in interstate commerce meet the official requirements of the U. S. P. and N. F. These activities require and, in fact, demand greater financial support from Congress than the FDA has received to date. It is inherently wrong for manufacturers to be required to pay for having their drugs certified as to purity and potency and, indeed, it has been just this program which brought about one of the developments for which the Food and Drug Administration merited its most severe criticism.

We have in the United States an ideal system whereby the standards of purity and strength of all important drugs are established by recognized professional experts working in behalf of the *United States Pharmacopeia* and *National Formulary*. These same standards then are used by the Food and Drug Administration in determining whether drugs in interstate commerce are suitable for use. This system has proved workable and effective and it should not be abandoned in favor of some new system which would prove both exceedingly expensive and open to all sorts of abuses not in the ultimate interest of the industry, the government, or the public which both serve.

L. F. TICE

STABILITY PROBLEMS WITH SOME VITAMINS IN PHARMACEUTICALS * †

By Thomas J. Macek ††

Vitamin and vitamin-mineral compositions, unlike other physiologically-active materials, are prepared in dosage forms primarily as mixtures. Such mixtures represent one of the largest, economically-important groups of modern pharmaceutical products. Interactions between vitamins and between the vitamins and other components may be found in such products, however, even though the vitamins are stable by themselves.

The literature and some personal experiences are reviewed. The discussion first summarizes the important decomposition pathways for vitamin A, thiamine, riboflavin, ascorbic acid, vitamin B₁₂, and pantothenic acid. The manuscript then considers interactions between thiamine and pantothenic acid, thiamine and cyanocobalamin, thiamine and riboflavin, cyanocobalamin and ascorbic acid, niacinamide and ascorbic acid and cyanocobalamin with folic acid.

VITAMIN and vitamin-mineral compositions, unlike many other physiologically-active materials, are prepared in dosage forms primarily as mixtures. Such mixtures represent one of the largest, economically-important groups of modern pharmaceutical products. Not unexpectedly, however, these multi-component products exhibit numerous problems with stability even though most of the individual vitamins are stable and may be prepared into reasonably stable dosage forms by themselves. Furthermore, certain of the decomposition reactions in multivitamin systems are not yet fully understood. This

* Presented at the Second Annual National Industrial Pharmaceutical Research Conference, Land O'Lakes, Wisconsin, June 12-15, 1960.

† A contribution from the Merck Sharp & Dohme Research Laboratories, West Point, Pennsylvania.

†† Director, Pharmaceutical Research and Development, Merck Sharp & Dohme, West Point, Pennsylvania.

prevails despite considerable progress in the art of preparing formulations having sufficient balance in potency to permit wide-scale, commercial distribution. In recent times, however, even this has been questioned, as in Canada, leading to various proposals for labeling products with expiration datings.

Interaction of vitamins may be found in all conventional formulations, including dry-filled and oil-filled gelatin capsules, compressed tablets, as well as in liquid oral and liquid parenteral products. At times, even inert formulation additives and diluents may be involved, particularly as they alter pH, supply moisture, or introduce trace metals or other reactive contaminants.


From the point of view of formulation, certain interactions of vitamins are more important than others. These will be discussed. Some prominent incompatibilities between the vitamins and non-vitamin formulation additives will also be noted. A full review of the very large literature on vitamins, however, is beyond the scope of this paper.

The more labile vitamin compounds are vitamin A, thiamine, riboflavin, ascorbic acid, vitamin B₁₂, and pantothenic acid. Discussion of the general problems of stability of each of these would seem worthwhile before proceeding with a consideration of some of the more specific interactions observed in product formulation.

Vitamin A—This compound occurs naturally in the animal organism as the free alcohol but predominantly in esterified form. For pharmaceutical use, it is today also prepared by synthesis. Esters such as the acetate and palmitate generally are preferred to the alcohol because of greater stability. The pure, crystalline acetate has been established as the international standard. Ethers have been made although some are biologically inactive. Vitamin A aldehyde possesses a biological potency equivalent to vitamin A alcohol, but is very unstable. Vitamin A acid has been obtained as a crystalline solid having about two-thirds of the activity of the alcohol. The properties of a number of derivatives are summarized in Table 1.

Vitamin A is sensitive to oxidation and readily undergoes autooxidation and polymerization. The oxidation is catalyzed by traces of metals, notably copper and iron, but the process is inhibited appreciably by the addition of antioxidants such as propyl gallate, hydroquinone, alpha-tocopherol and especially by combinations of

TABLE I
VITAMIN A₁ AND DERIVATIVES



	M.W.	M.P.	E 1% + 1 cm.	Biopotency I.U./Gm.
Alcohol	286.4	64°	1835	3.33 x 10 ⁶
Aldehyde	284.4	68°	1510	3.3
Acid	300.4	180°	1600	2.2

ESTERS

Acetate	328.5	57-8°	1550	2.91
Butyrate	356.5	---	1345	2.20
Laurate	468.7	---	1035	1.70
Palmitate	524.8	28-9°	975	1.60
Stearate	552.9	---	940	1.50
Benzoate	390.5	---	1240	1.80
Succinate	654.9	77-8°	1480	2.50

ETHERS

Methyl	300.5	34-5°	1660	3.50
Phenyl	362.5	90-2°	1460	0.10

+ λ_{Max}. 324-28 millimicrons (ethanol)

these antioxidants. High potency oil solutions of vitamin A esters containing antioxidants are used in liquid formulations and soft gelatin capsules. The acetate coated with gelatin or a gelatin-starch composition to give a stable, dry form is widely employed in dry capsules and tablets.

Vitamin A dissolved in vegetable oil can be destroyed when the oil becomes rancid owing to the formation of peroxides. The amount of decomposition varies with the kind of oil or fatty acid ester present but may be inhibited by the addition of an antioxidant, as indicated in Table 2.

TABLE 2
LOSS AFTER 150 HOURS AERATION *

		0.05%
	No Antioxidant	Propyl Gallate
Peanut Oil	76%	31
Ethyl Oleate	74%	31
Ethyl Stearate	47%	31
Mineral Oil	72%	45
Mineral Oil		
Ethyl Oleate	79%	36
Mineral Oil		
Ethyl Stearate	46%	25
Mineral Oil		
Glyceryl Monolaurate	69%	13

* 4 ml./sec. dry air passed through 750 IU vitamin A in 25 ml. solvent at 28-9° C.

Ref.—*Chem. Abst.*, 44, 4098 (1950).

Although thermal decomposition has been observed, vitamin A is quite stable when heated at moderate temperatures in an inert atmosphere in the dark. It is easily destroyed upon exposure to ultraviolet light, however, losing the typical absorption band at 325-328 millimicrons, forming biologically inactive products.

Vitamin A contains a series of five conjugated double bonds, four of which are in the side chain. Addition reactions, therefore, can occur with substances like bisulfite and certain solvents. With such high degree of unsaturation, numerous *cis-trans* isomers also

are possible (1). Natural vitamin A is all *trans*. This spatial arrangement is the most active biologically. The conversion of some of the double bonds into *cis* structures may decrease biological potency very appreciably although chemical assay may not be affected significantly. The properties of some geometric isomers are summarized in Table 3.

TABLE 3
PROPERTIES OF GEOMETRIC ISOMERS OF VITAMIN A PALMITATE

Isomer	USP XV Potency	Biopotency	Biopotency
	u/Gm.	u/Gm.	
All- <i>trans</i>	1.8×10^6	1.8×10^6	100
2-Mono- <i>cis</i>	1.30	1.37	105
6-Mono- <i>cis</i>	1.26	0.41	33
2,6-di- <i>cis</i>	1.32	0.41	31


Such changes have been reported recently to occur to the extent of about 20% of the vitamin A content in aqueous multivitamin dispersions and were readily detected by the reaction with maleic anhydride. The all *trans* and 6-mono-*cis* forms react rapidly to give an adduct which fails to give the typical blue color assay with antimony trichloride. The 2-mono-*cis* and 2,6-di-*cis* forms react slowly with maleic anhydride and thus can be distinguished from the more reactive forms. Infrared studies have been employed to support these findings.

Mineral acids destroy vitamin A and its geometric isomers and form anhydrovitamin A, a hydrocarbon practically devoid of biological activity. This decomposition involving the loss of a mole of water, as shown in Figure 1, was first obtained by the action of hydrogen chloride in ethanol (2). The kinetics of the reaction were studied more recently by Higuchi and Reinstein (3) who reported a significant rate when vitamin A acetate was dissolved in alcoholic or hydroalcoholic solutions.

Thiamine—Thiamine is found naturally as the free compound, as salts, as protein complexes, and as the ester with pyrophosphoric acid (cocarboxylase). Thiamine Hydrochloride U. S. P. is commercially synthesized as thiamine chloride hydrochloride and crys-

tallizes from aqueous-alcoholic solutions as the hemihydrate, which melts with decomposition at 248-250°C. The molar proportion of water is not constant during storage and may vary depending upon humidity conditions. A moisture content not exceeding 5% is permitted by the U. S. P.

FIGURE 1



In spite of its variable moisture, crystalline thiamine hydrochloride as a solid is stable for extended periods at room temperature and at elevated temperatures even in contact with air. Aqueous solutions of thiamine hydrochloride in the pH range 2.5-4.5 are stable for a considerable time.

Thiamine mononitrate, which crystallizes readily from water, is an anhydrous, non-hygroscopic form of the vitamin. It is soluble in water to the extent of 2.7 Gm. per 100 Gm. of water; the pH of such a solution is 6.8-7.1. Addition of a strong acid, e.g., HCl or nitric acid, to solutions of thiamine mononitrate increases solubility by virtue of the ionic equilibria involved. The addition of HCl to maintain a pH of 4, for example, results in concentrations of 18.5 Gm. thiamine mononitrate per 100 ml. at room temperature. The addition of a strong acid results in neutralization of the free amino group, the acidic pK of which is known to be 4.8.

Thiamine mononitrate does not deliquesce even on exposure to 90% relative humidity, differing markedly in this respect from thiamine hydrochloride and, therefore, is a more stable form of thiamine, particularly for dry or oil-based pharmaceutical preparations.

Solutions of thiamine at pH 3.5 can be heated at 120°C. without significant decomposition. The molecule decomposes appreciably above pH 5 and is extremely sensitive to heat in the neutral and alkaline pH range. Decomposition also occurs quite rapidly when alkaline solutions are merely stored at room temperature. There is increasing evidence that decomposition by hydrolysis may be a complex process yielding a number of products, the pyrimidine and thiazole moieties among them. Neither of these is biologically active.

In solution, pH appears to be the chief factor in the hydrolytic decomposition of thiamine although the literature contains numerous reports attributing stabilization and decomposition, both, to specific ions.

Thiamine is sensitive to both oxidation and reduction. Even mild conditions, such as storage of an alcoholic solution at room temperature for a long period of time, produce thiochrome by oxidation. This change occurs more rapidly as pH approaches 7. Since thiochrome possesses an intense blue fluorescence, it serves as a basis for assay following chemical oxidation of thiamine with alkaline potassium ferricyanide. In alkaline solutions, thiochrome is sensitive to light and the fluorescence disappears irreversibly. Thiamine can

be oxidized to a disulfide under physiological conditions at pH 7.5 with hydrogen peroxide or with oxygen. The same oxidation can be brought about in alkaline solution with iodine. The disulfide is fully active biologically. Mild reduction of thiamine with sodium hydrosulfide causes addition of a mole of hydrogen in the pyrimidine ring with almost complete loss of vitamin activity. Sodium sulfite, on the other hand, even at room temperature causes the so-called "sulfite cleavage" of the thiazole and pyrimidine portions, with nucleophilic addition of the sulfite to form an insoluble, biologically inert amino sulfonic acid.

For a long time, the industry was plagued by formation of trace, feather-like precipitates and discoloration in parenteral solutions of thiamine, particularly when exposed to light or heat. This problem today has largely been eliminated by the production of thiamine which is virtually metal-free, by proper choice of container components and by the discovery of numerous suitable stabilizers especially such compounds as monothioglycerol and thiosorbitol, some of which are covered by patents (4).

Riboflavin—Although widely distributed in nature, practically all of the riboflavin used in pharmaceuticals is prepared synthetically. The pure crystalline compound is only slightly soluble in water (12 mg. in 100 ml. at 27.5°C.). Because of its limited solubility and the desire to obtain more concentrated pharmaceutical solutions, riboflavin often has been combined with other compounds capable of increasing its solubility. These include N-methylacetamide, L-tyrosine amide, tryptophan, sodium acetyl tryptophan, urea, niacinamide, urethane, veratryl alcohol, gallic acid, salts of boric acid, propylene glycol, benzyl alcohol, and sodium 3-hydroxy-2-naphthoate. Most of these solubilizers have been described in patents. The most useful for pharmaceuticals probably are propylene glycol and niacinamide. At pH 5, riboflavin solubility increases approximately 250 times when niacinamide concentration is increased from 5 to 50 per cent. A super-saturated solution of riboflavin alone or with added solubilizer may be stable physically, often requiring days to crystallize. Some crystalline forms of riboflavin incidentally are alleged to be more rapidly soluble in water than others. Riboflavin also can be fused with solubilizing compounds such as urea, urethane, or niacinamide to give products which yield aqueous solutions up to 6% riboflavin concentration (5).

Water-soluble chemical derivatives include esters with succinic, citric, and other organic acids. However, not all of these are fully active biologically. On the other hand, the 5'-phosphoric acid ester is quite soluble as the sodium salt, is fully active biologically, and is an item of commerce. It possesses the incompatibilities of the phosphate ion, however, precipitating with calcium, with iron and other cations.

Crystalline riboflavin is stable at ordinary temperatures but exposure to light slowly destroys vitamin activity. In solution, stability of riboflavin is influenced largely by pH. Acid or even neutral aqueous solutions are stable to heat if protected from light and can be sterilized by autoclaving. Alkali decomposes riboflavin rapidly.

The rate of destruction by light becomes greater with increasing temperature and pH. Photolysis of riboflavin in neutral solution in the absence of oxygen causes disappearance of the color with the formation of deuterio-leuco-riboflavin. This compound can be dehydrogenated by oxygen. Alkali subsequently converts the dehydrogenated compound to lumiflavin. Photolysis in alkaline solution also produces lumiflavin, whereas photolysis in acid or neutral solution produces lumichrome, an intensely fluorescent compound.


On reduction even with such mild agents as sodium thiosulfate, riboflavin readily takes up two hydrogen atoms. The dihydrocompound is colorless, shows no fluorescence, and is called leuco-riboflavin. It is readily oxidized to riboflavin by air. Actually, three crystalline compounds can be obtained by stepwise reduction or hydrogenation of riboflavin, but these are of little consequence in pharmaceuticals.

Ascorbic Acid—1-Ascorbic Acid, a white crystalline optically-active solid, which melts with decomposition at 190-2°C., is a strong monobasic acid. It liberates carbon dioxide from carbonates and bicarbonates, and forms well-defined salts with cations such as sodium, potassium, calcium, and magnesium.

The outstanding feature of the structure is the presence of an α -keto-diol system involving carbons 1, 2 and 3 as shown in Figure 2. This portion of the molecule is responsible for the characteristic ultra-violet absorption, for the high acidity of the compound, and also for the well-known reducing properties of 1-ascorbic acid.

FIGURE 2

ASCORBIC ACID



In solution at pH 2, there is an intense band of absorption in the ultraviolet with a maximum at 244 millimicrons and an extinction coefficient ($A_{1\text{ cm}}^{1\%}$) of approximately 500. The absorption changes gradually as pH increases above pH 2. At pH 6 to pH 10, the solutions have a maximum at 266 millimicrons; more strongly alkaline solutions have a maximum at approximately 294 millimicrons.

A 1% solution of ascorbic acid has a pH approximately 2.5. Titration of this solution to pH 8 requires one equivalent of alkali. Although both hydroxyls of the ene-diol system are acidic, the first dissociation constant ($pK_1 = 4.1$) is associated with the hydroxyl at carbon 3. The second dissociation constant ($pK_2 = 11.6$) is associated with the hydroxyl at carbon 2.

Like all α -keto-ene-diols, 1-ascorbic acid is a powerful reducing agent in acid and neutral solutions. It reduces Fehling's solution and ammoniacal silver nitrate, and reacts with iodine and other halogens and with dichlorophenol-indolphenol which is employed in its analysis. It also possesses chemical properties of a typical carbohydrate molecule, forming a bisphenylhydrazone with two moles of phenylhydrazine. The lactone portion of the molecule is remarkably resistant to hydrolysis. The principal decomposition of ascorbic acid, therefore, is oxidation, to which it is very sensitive, especially in aqueous

solution. The rate of oxidation increases in the presence of metals, especially copper or iron.

The first oxidation product of ascorbic acid is dehydroascorbic acid. This reaction is reversible and dehydroascorbic acid is reduced to ascorbic with hydrogen sulfide or HI. 1-Dehydroascorbic acid possesses the full biological activity of 1-ascorbic acid and, indeed, some of the naturally-occurring vitamin exists in the oxidized form. Further oxidation of 1-dehydroascorbic acid, however, involves extensive degradation to oxalic acid and 1-threonic acid and loss of activity. This decomposition is accelerated first by air and then by alkali. Neutral solutions of sodium ascorbate thus can be made in the absence of air which retain potency for long periods of time. Oxidation of ascorbic acid in vehicles such as glycerin, propylene glycol, sorbitol, or concentrated sugar solutions is low as shown by the data in Table 4. Stability of 1-ascorbic acid improves in aqueous

TABLE 4
ASCORBIC ACID
STABILITY IN VARIOUS VEHICLES

<i>Conc. = 100 mg./ml.</i>				
	<i>Water</i>	<i>Syrup USP</i>	<i>90% Glycerin</i>	<i>Propylene Glycol</i>
Initial	100.0%	100.0%	100.0%	100.0%
RT-1 month	92.3%	98.0%	98.5%	97.0%
RT-3 months	79.0%	95.8%	97.0%	93.5%
RT-6 months	58.0%	86.5%	90.8%	85.7%
RT-12 months	34.7%	80.3%	89.0%	82.0%
<i>Conc. = 5 mg./ml.</i>				
	<i>Water</i>	<i>Corn Syrup</i>	<i>Sorbitol</i>	<i>3% Methocel</i>
Initial	100.0%	100.0%	100.0%	100.0%
RT-1 month	90.0%	96.5%	99.0%	76.0%
RT-3 months	74.5%	92.0%	99.0%	39.5%
RT-6 months	40.5%	90.0%	96.0%	—
RT-12 months	—	76.0%	89.0%	—

Data from Bandelin & Tuschhoff, *J. A. Ph. A.* 44, 241 (1955).

solution with increasing concentrations of sodium chloride as shown in Table 5. The improvement in stability in these vehicles generally is associated with a lower concentration of dissolved oxygen in the vehicle. Addition of gums merely to increase viscosity of aqueous solutions is insufficient to exert an equivalent stabilizing effect.

TABLE 5
ASCORBIC ACID
STABILITY IN SALT SOLUTION

	Water	Conc. = 1 mg./ml.—% Retained		
		1% NaCl	2% NaCl	5% NaCl
6 Hours	77%	97%	100%	98%
24 Hours	55%	87%	97%	99%
3 Days	43%	63%	88%	99%
5 Days	5%	16%	65%	91%
7 Days	—	—	52%	93%
9 Days	—	—	39%	86%

Vitamin B₁₂—Cyanocobalamin, the red crystalline form of vitamin B₁₂, was first isolated in pure state in 1948. In aqueous solution, this compound has a characteristic absorption spectrum with peaks at 278, 361, and 550 millimicrons. The spectrum is not markedly affected by small changes in pH.

Early investigations of structure revealed the presence of cobalt, phosphorus, and the cyano group. Potentiometric titration showed cyanocobalamin to be a polyacidic base containing six weakly basic groups. Further studies yielded evidence of a complex organic moiety which could be degraded to yield 5,6-dimethylbenzimidazoles and ribazoles, the ribofuranisido substituents of the former.

But despite its large organic molecular composition, cyanocobalamin essentially is a coordination compound of cobalt. Its chemistry and stability are best viewed in terms of the properties of this inorganic coordination complex. The cyano group, though very weakly ionized, is replaceable by other ions or molecules. The number of B₁₂ modifications is limited only by the number of ions or molecules which can replace cyanide in the cobalt coordination complex. In the formation of vitamin B_{12a}, or hydroxocobalamin, the


cyanide ion is liberated and replaced by a hydroxo group. The cyano group may also be replaced by chloride, nitrate, acetate, and other anions to form the respective B₁₂ analogs. Some of the latter are almost completely ionized. The general equilibrium for these transformations is represented in Figure 3.

FIGURE 3



Cobalamin

Cobalaminium Ion




Hydroxocobalamin
(Cobalaminium Hydroxide)

Symbols

- | | | |
|----------------------|---|--|
| - = negative charges | } | - To satisfy the remaining coordination bonds. |
| o = electron pairs | | |

Aqueous solutions of cyanocobalamin are most stable in the pH range of 4 to 7 at normal temperatures. Exposure to sunlight brings about loss of microbial activity. Autoclaving at pH 4.5 causes a slight decomposition but is probably of no consequence in pharmaceuticals. The presence of reducing agents, however, even in trace quantities, frequently introduced by way of formulation additives, can markedly influence potency as a result of decomposition of the reduced form to irreversible oxidation products. This is shown diagrammatically in Figure 4.

FIGURE 4



The outstanding incompatibility of cyanocobalamin, of course, is that with ascorbic acid. Decomposition products of thiamine, ferrous salts, sodium bisulfite, reducing sugars, and some flavors are equally destructive, however. Some of these will be considered presently.

Pantothenic Acid—d-Pantothenic acid itself is a yellow viscous oil and not used in pharmacy. It forms a crystalline water-soluble calcium salt, however, which is widely employed. The alcohol analog of pantothenic acid also is available in two forms. d-Pantotheryl alcohol is a viscous, colorless liquid and dl-Pantotheryl alcohol, a crystalline solid. Both forms are soluble in water and alcohol but not in oil. The dl-form is half as active biologically.

All forms of this vitamin are sensitive toward acids and bases, and to heat. The range of greatest stability is pH 4 to 7 with optimum stability at pH 6. Hydrolysis occurs at the amide linkage between the lactone and the β -alanine moieties and proceeds at an increasing rate as the pH varies from the optimum. The compound has high structural specificity so that hydrolysis is accompanied by loss of biological activity.

Thiamine and Pantothenic Acid—The kinetics of the hydrolysis of thiamine and pantothenate in acid solution was reported in 1944 by Frost and McIntire (6). The hydrolysis of pantothenate is a first-order reaction with respect to pantothenate concentration. The decomposition of thiamine, on the other hand, is much more complicated yielding several decomposition products as pH varies as shown by the paper chromatograms of Busse *et al.* (7).

The opposing stability relationships between thiamine and pantothenate as regards pH present practical difficulties in formulation, especially of liquid multivitamin products. To achieve optimum stability of thiamine, riboflavin, ascorbate, pyridoxine, and even of cyanocobalamin, multivitamin liquids usually are prepared at about pH 4. Under these circumstances, pantothenate is rapidly decomposed, potency often falling below label claim on storage of products even with the addition of large manufacturing overages.

The development of pantothenyl alcohol as a more stable analog of pantothenic acid for use in liquids aided in the stability problem. The rate of decomposition of pantothenyl alcohol was significantly less than pantothenate at pH 4 in both pure aqueous solution and in liquid compositions with other vitamins. In the more stable pH range of 6 to 8, however, the rates of hydrolysis of pantothenyl alcohol and calcium pantothenate were essentially the same (8).

The stability of the amide linkage in pantothenic acid to hydrolysis is influenced by the polarity of the terminal groupings on the molecule. The hydroxyl of pantothenyl alcohol is less polar than the terminal carboxyl of pantothenic acid. The inductive effect of the terminal carboxyl group increases the positive charge on the nitrogen and weakens the resonance of the amide. This is especially noted in mild acid solution. Weakening of the amide resonance favors hydrolysis (9).

In dry B complex mixtures, it has sometimes been the practice to add an acidic substance, such as tartaric acid, to lower pH and aid in the stability of thiamine HCl. Such a practice obviously is unfavorable to the stability of calcium pantothenate. When thiamine mononitrate is employed, the addition of an acidic stabilizer becomes unnecessary (10). Thiamine mononitrate, therefore, is much more compatible with calcium pantothenate in dry mixtures intended for use in capsules and tablets. Data showing differences in stability are shown in Table 6.

TABLE 6
DRY-FILLED B VITAMIN CAPSULES

	A	B	C
	<i>Thiamine (3 mg./Capsule)</i>		
Initial	100%	100%	100%
1 mo.-RT	62%	95%	100%
6 mo.-RT	17%	93%	100%
1 mo.-40°	32%	97%	95%
3 mo.-40°	21%	95%	97%
	<i>Calcium Pantothenate (10 mg./Capsule)</i>		
Initial	100%	100%	100%
1 mo.-RT	100%	92%	100%
6 mo.-RT	100%	65%	100%
1 mo.-40°	100%	62%	99%
3 mo.-40°	100%	16%	96%

A—Thiamine HCl, no tartaric acid.

B—Thiamine HCl with tartaric acid.

C—Thiamine Mononitrate, no tartaric acid.

In oil suspension compositions such as used in filling soft gelatin capsules, the stability of calcium pantothenate is improved when dibasic magnesium phosphate is added as a stabilizing component (11).

Thiamine and Cyanocobalamin—In one of the first published reports on stability studies with cyanocobalamin, Macek and Feller (12) observed a loss of 16% of cyanocobalamin ($C = 15$ mg./ml.) upon autoclaving an acetate buffered solution at pH 4 containing a large excess of thiamine hydrochloride ($C = 100$ mg./ml.). However, negligible loss of cyanocobalamin was observed when this same solution was stored at room temperature for a period of one year. It was concluded that there was no incompatibility between thiamine and cyanocobalamin.

Shortly thereafter, Blitz *et al.* (13) reported that cyanocobalamin was not stable in a B-complex parenteral solution at pH 4.25 containing both thiamine and niacinamide among its constituents. The large losses of cyanocobalamin observed by these workers could not be reproduced in subsequent experiments by the earlier workers. The B vitamin solutions were prepared at pH 3.0 and 4.5 and were steri-

lized without heat. They were stable even after prolonged storage (14). The discrepancy between the two studies was resolved, however, when it was realized that Blitz *et al.* employed concentrations of both thiamine and niacinamide that were substantially greater. Furthermore, the decomposition was easily accelerated by heat, or by the presence of thiamine decomposition products, notably the thiazole moiety, which functioned as a reducing agent for cyanocobalamin (15). The pyrimidine moiety had no effect on cyanocobalamin stability, nor did thiochrome (16).

More recent studies by Indian workers (17, 18) have confirmed these findings but have also proposed that the deterioration could be prevented by the use of iron salts. These studies indicated that the same iodine-titratable reducing substances were formed by the thermal decomposition of thiamine at pH 4.0-4.5 in the presence of niacinamide, as when a solution of thiamine alone decomposed at pH 8. Both solutions containing these reducing substances, which indeed may have been the same, were capable of destroying cyanocobalamin. Ferric chloride added to the thiamine-niacinamide solution at pH 4.0-4.5 spared cyanocobalamin from reduction by the decomposition products. Stabilization of cyanocobalamin by ferric chloride also was noted in solutions prepared at pH 8 but was incomplete. A summary of these findings is given in Table 7.

TABLE 7

STABILIZATION OF VITAMIN B₁₂ BY FeCl₃

pH	Niacinamide	FeCl ₃	Potency After Test	
	mg./ml.	mg./ml.	B ₁ (mg./ml.)	B ₁₂ (mcg./ml.)
4.0	—	—	11.0	4.5
8.0	—	—	7.5	0.5
8.0	—	0.5	6.2	2.5
4.0	100	—	10.6	nil
4.0	100	0.5	10.6	4.5
8.0	100	—	7.0	0.5
8.0	100	0.5	5.1	2.5

Ref.—Mukherjee & Sen, *J. Pharm. Pharmacol.* 11, 26 (1959).

Initial Conc.—Thiamine = 15 mg./ml.

Vitamin B₁₂ = 5 mcg./ml.

Test—Heat at 100°C./4 hours.

The observations of the effect of thiamine and niacinamide on cyanocobalamin also were confirmed by Gambier and Rahn in Brazil (19). But these workers defined a critical ratio, a critical pH, and special conditions for processing the parenteral solutions which excluded heat. Stable solutions containing thiamine, niacinamide, and cyanocobalamin could be prepared by these special means if the thiamine-cyanocobalamin ratio did not exceed 120 to 1. If the thiamine-cyanocobalamin ratio was 5,000 to 1 or 10,000 to 1, then even the refined techniques failed to prevent decomposition of cyanocobalamin.

The problem of cyanocobalamin decomposition by thiamine is especially acute in high-potency parenteral products. It is much less likely to occur in maintenance-type oral liquids where concentrations invariably are smaller and where non-aqueous vehicle components contribute toward stabilization. Nevertheless, methods have been devised for minimizing the incompatibility in parenteral solutions and stable products are available commercially. One method which assures maximum stabilization involves lyophilization.

Thiamine and Riboflavin—An incompatibility between thiamine and riboflavin has been reported in aqueous B-complex solutions which manifests itself in physical changes involving trace precipitation of thiochrome or chloroflavin (20).

The precipitation of thiochrome is induced by the oxidative action of riboflavin on thiamine. It occurs more rapidly, the greater the riboflavin concentration. The presence of a greater quantity of air in the ampuls over the solutions also accelerates the process. On the other hand, the reducing action of thiamine or its decomposition products on riboflavin is denoted by the crystallization of chloroflavin. It occurs more readily as the ratio between thiamine and riboflavin concentrations increases. When the chloroflavin crystals in decomposed solutions are redissolved, full recovery of riboflavin potency is attained. The incompatibility has been reported to be completely eliminated in solutions containing thiamine, riboflavin, pyridoxine, and niacinamide by the addition of ascorbic acid (21).

Cyanocobalamin and Ascorbic Acid—The marked decomposition of vitamin B₁₂ and vitamin B₁₂ concentrates in aqueous solutions of ascorbate was first reported in 1949 by Gakenheimer and Feller (22). Subsequent studies showed that the rate of loss of cyanocobalamin in

ascorbate solution at pH 2.5 and 3.0 was approximately 1.5 per cent per day at room temperature. Vitamin B_{12a}, on the other hand, when examined under identical conditions, showed practically complete loss of color and activity within one day. The cyanocobalamin-ascorbate reaction was observed to progress faster at higher pH and was complete in less than one hour at pH 7 (23).

In another study, ascorbic acid was reported to be destructive for non-cyanocobalamins such as chloro-, nitro- and thiocyanatocobalamin, decomposition in pH 4 buffer requiring only a few hours at 25°C. Furthermore, the stability of vitamin B₁₂ concentrates to ascorbate varied directly with their cyanocobalamin content (24).

In view of this prominent incompatibility and the desire on the part of manufacturers and clinicians alike to have both vitamins present in the same multivitamin formulation, a great deal of work has been done in recent years with an aim toward stabilizing the B₁₂-ascorbate reaction. Many totally-unrelated methods were developed for improving the stability of cyanocobalamin in the ascorbate system and in other compositions. These are generally described in patents. A brief review of some of these will be interesting.

Soluble, non-toxic bitartrate salts at a preferred concentration of 0.1% have been employed to stabilize vitamin B₁₂ concentrates against destruction by heat (25). Thiodipropionic acid (0.1%) was also described for the same purpose (26). Reasonably stable solution compositions containing crystalline vitamin B₁₂, or a mixture of B₁₂ and B_{12a} derived from concentrates, with ascorbic acid and thiamine were claimed to be produced by cooperating stabilizers which included Tween 20, cysteine HCl and sodium copper chlorophylline or the sodium salt of yeast nucleic acid (27). Ammonium sulfate (100-10,000 mcg. per ml.) has been employed to prepare storage-stable solutions of vitamin B₁₂ concentrates which contained substantial proportions of hydroxocobalamin (28). To overcome destruction of B₁₂ by ferrous salts, a composition containing ferric versenate was described (29). Resin adsorbates of vitamin B₁₂ on cation exchangers were claimed to have unusual stability, especially under acid conditions in the presence of ascorbic acid, ferrous iron, and other reducing agents (30). More recently, α -hydroxynitriles (mandelonitrile) (31), non-toxic cationic salts of metalocyanide complexes (potassium ferrocyanide, potassium cobalticyanide) (32), non-toxic nitrites (33), and soluble molybdenum compounds (sodium or ammonium molyb-

date) (34), all have been claimed as stabilizing agents for the B_{12} -ascorbate decomposition.

However, perhaps the most significant method for improving the stability of these very troublesome cyanocobalamin systems arose from the observation of Skeggs (35), who found that the activity of aqueous solutions of vitamin B_{12} concentrates could be improved considerably by the addition of saccharated iron oxide. The idea of using soluble iron salts was successfully applied to the stabilization of aqueous solutions of cyanocobalamin and ascorbic acid by Newmark (36), wherein the non-toxic iron compound was added in the range of 17-1700 mcg. per ml. This low concentration of iron, surprisingly, did not appear to affect ascorbic acid stability and in many cases did, indeed, stabilize the cyanocobalamin component. Some data showing the stabilizing effect of saccharated iron oxide on the cyanocobalamin-ascorbate system are given in Table 8. This system now is used rather widely commercially for producing oral, liquid multivitamin products.

TABLE 8

STABILITY OF CYANOCOBALAMIN AND ASCORBIC ACID IN LIQUIDS
WITH AND WITHOUT SACCHARATED IRON OXIDE (50 mcg./ml.)

	<i>With Fe</i>		<i>No Fe</i>	
	B_{12}	C	B_{12}	C
		<i>Polyvitamin Drops</i>		
Initial	100%	100%	100%	100%
50°C.-20 days	61%	62%	Trace	31%
37°C.- 1 month	100%	90%	Trace	97%
RT - 2 months	92%	95%	30%	97%
RT - 4 months	92%	89%	—	—
		<i>Polyvitamin Syrup</i>		
Initial	100%	100%	100%	100%
50°C.-10 days	92%	92%	Trace	80%
50°C.-30 days	59%	79%	Trace	45%
37°C.- 1 month	82%	93%	69%	96%
RT - 4 months	—	—	60%	94%

Another process for stabilizing the cyanocobalamin-ascorbate reaction employs a water-swellable clay of the Montmorillonite type (Veegum, Thix) (37).

Preliminary studies by Bartilucci suggested that the decomposition products of ascorbic acid were involved in the destruction of cyanocobalamin in aqueous solution (38). Later work provided evidence that increasing concentrations of dehydroascorbic acid produced progressively greater losses of cyanocobalamin in ascorbic acid solution (39).

Finally, commercial 70% sorbitol solution has been employed as a vehicle, alone or in mixture with glycerin, for oral liquid products containing ascorbic acid, cyanocobalamin, and ferrous gluconate. Circumvention of the mutual incompatibilities in such preparations even after prolonged storage at room temperature was attributed largely to the binding of water by sorbitol. The possible complexation between sorbitol and either or both ascorbic acid and cyanocobalamin also was suggested (40).

The studies of Barr *et al.* (41) showed that cyanocobalamin alone was stable in sorbitol and in glycerin, whereas dextrose and sucrose solutions caused decomposition during storage at 25°C. and 45°C. In this connection, Bartilucci (42) cautioned that sorbitol containing trace metal contaminants as a result of improper storage was destructive to ascorbic acid.

Miscellaneous—Two other interactions between vitamins are worth noting. The first is the canary-yellow "addition compound" that forms when equimolar quantities of ascorbic acid and niacinamide are mixed, especially in the dry state. This reaction product maintains essentially the full potency of each constituent but has a different melting point than either component. A patent issued to Fox and Opferman (43) teaches that, unless this compound is preformed, subsequent difficulty may arise with thickening and hardening of the mixtures employed in filling soft gelatin capsules.

The second interaction involves the incompatibility of folic acid in cyanocobalamin solutions adjusted to pH 6. At this pH, 5 mg. per ml. of folic acid is stable and remains in solution only if niacinamide also is added to maintain clarity and serve as a stabilizing component (44).

Finally, this paper would be incomplete without reference to the excellent studies of Garrett dealing with the prediction of stability of multivitamin products (45).

A great deal more could be said about methods of formulating multivitamin products, choice of granulating methods and coating

procedures in tablet manufacture, selection of solubilizers and vehicles for liquids, vitamin-mineral combinations, and many other topics. Despite the numerous problems, various ways have been devised for producing multivitamin and vitamin-containing combinations having acceptable stability in both solid and liquid form. Much of this is due to the ingenuity and resourcefulness of research pharmacists in searching out and developing techniques, solvents, derivatives, and stabilizing additives. Their efforts have contributed substantially toward the establishment of numerous stable and useful products currently available in world-wide markets.

REFERENCES

- (1) Zechmeister, *Chem. Revs.* 34, 267 (1944).
- (2) Edisbury, J. R., Gillam, A. E., Heilbron, I. M., and Morton, R. A., *Biochem. J.* 26, 1164 (1932).
- (3) Higuchi, T., and Reinstein, J. A., *J. A. Ph. A., Sci. Ed.* 48, 155 (1959).
- (4) Bray, M. D., U. S. Pat. 2,498,200.
- (5) U. S. Pat. 2,840,517 (8-30-49).
- (6) Frost, D. V., and McIntire, F. M., *J. Am. Chem. Soc.* 66, 425 (1944).
- (7) Lhoest, W. J., Busse, L. W., and Baumann, J. A. Ph. A., *Sci. Ed.* 47, 254 (1958).
- (8) Rubin, S. H., *ibid.* 37, 502 (1948).
- (9) Hammett, L. P., *Physical Organic Chemistry*, Ed. 1. New York: McGraw-Hill, 1940, page 365.
- (10) Macek, T. J., Feller, B. A., and Hanus, E. J., *J. A. Ph. A. Sci. Ed.* 39, 365 (1950).
- (11) Macek, T. J., U. S. 2,510,487, August 22, 1950.
- (12) Macek, T. J., and Feller, B. A., *J. A. Ph. A., Sci. Ed.* 41, 285 (1952).
- (13) Blitz, M., Eigen, E., and Gunsberg, E., *ibid.* 43, 651 (1954).
- (14) Macek, T. J., and Feller, B. A., *ibid.* 44, 662 (1955).
- (15) *Ibid.*, p. 254.
- (16) Ravin, L. J., and Doerge, R. F., *ibid.* 48, 425 (1959).
- (17) Mukherjee, S. L., and Sen, S. P., *J. Pharm. Pharmacol.* 9, 759 (1957).
- (18) *Ibid.* 11, 26 (1959).
- (19) Gambier, A. S., and Rahn, E. P. G., *J. A. Ph. A., Sci. Ed.* 47, 356 (1958).
- (20) Gambier, A. S., Jagle, B. S., and Rahn, E. P. G., *Ciencia e cultura* 6, 171 (1954).
- (21) Gambier, A. S., and Rahn, E. P. G., *J. A. Ph. A., Sci. Ed.* 46, 134 (1957).

- (22) Gakenheimer, W. C., and Feller, B. A., *ibid.* 38, 660 (1949).
- (23) Trenner, N. R., Buhs, R. P., Bacher, F. A., and Gakenheimer, W. C., *ibid.* 39, 361 (1950).
- (24) Hutchins, H. H., Cravioto, P. J., and Macek, T. J., *ibid.* 45, 806 (1956).
- (25) DeRose, A., U. S. 2,566,123, August 28, 1951.
- (26) Leffler, M. T., U. S. 2,579,679, October 25, 1951.
- (27) Winsten, W. A., U. S. 2,662,048, December 8, 1953.
- (28) Michel, G. H., and Knight, K. W., U. S. 2,778,771, January 22, 1957.
- (29) Sahyun, M., U. S. 2,804,423, August 27, 1957.
- (30) Bouchard, E. F., and Friedman, I. J., U. S. 2,830,933, April 15, 1958.
- (31) Conine, J. W., and Zuck, D. A., U. S. 2,835,627, May 20, 1958.
- (32) Zuck, D. A., U. S. 2,874,089, February 17, 1959.
- (33) MacDonald, L. H., U. S. 2,914,446, November 24, 1959.
- (34) Buchler, H. J., U. S. 2,923,663, February 2, 1960.
- (35) Skeggs, H., U. S. 2,584,627, February 5, 1952.
- (36) Newmark, H. L., U. S. 2,823,167, February 11, 1958.
- (37) Bryant, C. C., U. S. 2,846,352, August 5, 1958.
- (38) Bartilucci, A., and Foss, N. E., *J. A. Ph. A.*, Sci. Ed. 43, 159 (1954).
- (39) Bartilucci, A., DiGirolamo, R., and Eisen, H., *ibid.* 47, 42 (1958).
- (40) Gerber, C. F., Hetzel, C. P., Klioze, O., and Leyden, A. F., *ibid.* 46, 635 (1957).
- (41) Barr, M., Kohn, S. R., and Tice, L., *ibid.*, p. 650.
- (42) Bartilucci, A. J., DiGirolamo, A., and Eisen, H., *ibid.*, p. 627.
- (43) Fox, S. H., and Opferman, L. P., U. S. 2,433,688, December 30, 1947.
- (44) Taub, A., and Lieberman, H., *J. A. Ph. A.*, Sci. Ed. 42, 183 (1953).
- (45) Garrett, E. R., *ibid.* 45, 171 and 470 (1956).

BOOK REVIEW

Heterocyclic Chemistry. A. R. Katritzky and J. M. Lagowski.
274 pp. John Wiley & Sons, Inc., New York, N. Y., 1960.
Price: \$4.75.

The authors have endeavored to construct a text on heterocyclic chemistry which is intended to indicate to the student that brute memory is not a prerequisite to the comprehension of heterocyclic syntheses. The approach to reaction sequences is wholly electronic in nature and should appeal to the educator and student with a leaning toward the modern concepts of organic chemistry.

The subject matter covers three through eight membered ring systems, with the emphasis on five and six membered compounds. Each chapter is subdivided to consider reactions of the heterocyclic and non-heterocyclic ring systems in the case of fused ring systems. A short chapter of ring systems involving atoms other than nitrogen, sulfur, or oxygen is also included.

For the most part, the nomenclature follows the *Chemical Abstracts* system with only slight reversion to European nomenclature.

From the pedagogic viewpoint, the lack of review questions and the meagre treatment of physical data, to perhaps correlate with electronic mechanistic interpretations, detracts somewhat from the usefulness of the book as an undergraduate text. Also, no doubt in the desire to be complete, the text includes a tremendous wealth of information and some topics are treated sketchily, a source of confusion even for the more advanced student.

It is refreshing, however, to see this new and logical approach to heterocyclic organic chemistry, and the authors must be commended for their organization and treatment of the subject in a rational manner. The text is highly recommended for the student and research worker desirous of emerging from the "cook-book" status of heterocyclic chemistry.

A. R. GENNARO

INDEX TO VOLUME 132 OF THE AMERICAN JOURNAL OF PHARMACY

AUTHOR INDEX, 1960

(B)—Book Review

(E)—Editorial

	PAGE		PAGE
Baer, John E.—		Ehrenstein, Elsa—	
Evaluation of Diuretic Agents	5	Subsidia Pharmaceutica 1959	
Barr, Martin—		(B)	36
Vitamin A (B)	115	Subsidia Pharmaceutica and	
Bendoraitis, J. G.—		Index Nominum 1960 (B) .	304
Composition Studies on To-		Gennaro, A. R.—	
bacco	165	Heterocyclic Chemistry (B) ..	456
Beyer, Karl H.—		Lipide Chemistry (B)	423
Evaluation of Diuretic Agents	5	Organic Reactions, Vol. X	
Bhatnagar, J. K.—		(B)	423
Histological Studies of the		Organic Syntheses, Vol. 39	
Genus Lavandula	252	(B)	37
Buday, Paul V.—		Oxidation—Reduction Potentials	
Neurohormone - Phrenotropic		of Organic Systems	
Drug Activity Correlates ..	355	(B)	241
Burlage, Henry M.—		The Story of Chemistry (B)	347
Sequence of Amino-Acids in		Gershenfeld, Louis—	
Proteins and Polypeptides ..	282	The 1960 Remington Medalist	
Connor, John D.—		and His Co-Workers	428
Sequestration and Its Applica-		Gjerstad, Gunnar—	
tions	325	Gibberellic Acid	98
Cooper, Jack—		Gupta, V. D.—	
Pharmacist and Research, The	158	Sequence of Amino-Acids in	
Davis, William A.—		Proteins and Polypeptides ..	282
Evaluation of Drugs Affecting		Hall, Nathan A.—	
Neuro-Muscular Transmis-		A Note on the Solubility of	
sion	112	Salicylic Acid in Isopro-	
Dunn, Marin S.—		panol-Water Solutions at	
Histological Studies of the		25°C.	406
Genus Lavandula	252	Haynes, Morris—	
		Gibberellic Acid	98

	PAGE		PAGE
Keith, Eaden F.—		Sampey, John R.—	
Evaluation of Analgesic Sub-		Decade of Chemotherapeutic	
stances	202	Management of Chronic	
Kramer, John E.—		Granulocytic and Chronic	
Contribution to Western Penn-		Lymphocytic Leukemia	145
sylvania Pharmacy, A. (B)	37	Decade of Chemotherapeutic	
Medieval and Renaissance		Management of Hodgkin's	
Medicine (B)	38	Disease	27
Lee, Charles O.—		Decade of Management of	
Shakers as Pioneers in the		Polycythemia Through	
American Herb and Drug		Chemotherapy	338
Industry	178	New Nitrogen Mustards in the	
Macek, Thomas J.—		Management of Hodgkin's	
Stability Problems with Some		Disease	231
Vitamins in Pharmaceuticals	433	Phosphoramides in the Man-	
Madjor, A.—		agement of Lymphomas and	
Exponential De-Emulsification		Reticuloses	295
and Stability of Pharmaceu-		The Unique Place of Folic	
tical Emulsions	315	Acid Antagonists in Control-	
Menczel, E.—		ling Malignant Blood Dis-	
Exponential De-Emulsification		eases	415
and Stability of Pharmaceu-			
tical Emulsions	315	Smith, Thomas H. F.—	
Neville, Jere—		A Chronology of Atherosclero-	
Plurionics as a Suppository		sis	390
Base	301	An Introduction to Pharma-	
Osol, Arthur—		cology (B)	306
Methoden der Organischen		Biophysical Science: A Study	
Chemie (Houben-Weyl),		Program (B)	273
Vol. 1, Part 2 (B)	195	Connective Tissue, Thrombo-	
Methoden der Organischen		sis and Atherosclerosis (B)	39
Chemie (Houben-Weyl),		Fat Consumption and Coro-	
Vol. 5, Part 4 (B)	305	nary Disease: The Evolu-	
Rabinovitz, M.—		tionary Answer to the Prob-	
Exponential De-Emulsification		lem (B)	276
and Stability of Pharmaceu-		Figures that Count: Mathe-	
tical Emulsions	315	matics for Nurses (B)	275
Rossi, G. Victor—		New and Non-Official Drugs,	
Psychotherapeutic Drugs	86	1959 (B)	274
Rusaniwskyj, W.—		Nutrition and Atherosclerosis	
Composition Studies on To-		(B)	38
bacco	165	Stedman, R. L.—	
		Composition Studies on To-	
		bacco	165

	PAGE		PAGE
Swafford, William B.—		Project HOPE (E)	388
Carbopol 934 as Ephedrine Sul-		Unpublished Abstracts of Ar-	
fate Jelly Base	383	ticles on Pharmaceutical	
Pluronic as a Suppository		Subjects, Series 8 (B)	115
Base	301		
Swain, A. P.—		Vogin, Eugene E.—	
Composition Studies on To-		A Review of Bioflavonoids ...	363
bacco	165	Biochemical Preparations, Vol-	
		ume 7 (B)	386
		Drugs, Their Nature, Action	
		and Use (B)	78
Tice, L. F.—		Modern Pharmacology and	
A. M. A. and Proprietary		Therapeutics (B)	196
Drugs (E)	335	Principles of Biochemistry (B)	242
An Example for Others (E)	250	Principles of Human Pathol-	
Drug Counterfeiting (E)	313	ogy (B)	242
Drug or Beverage? (E)	84	The Neurochemistry of Nu-	
Drug Research Today (E) ...	44	cleotides and Amino Acids	
Expansion of Drug Certifica-		(B)	307
tion by the FDA (E)	431		
Generic Name Pitfall, The (E)	2		
Government Purchases of		Ware, Ralph M., Jr.—	
Drugs Overseas (E)	280	Some Objectives of the	
Need for Mutual Support (E)	156	N. A. B. P.	409
Need for Professional Orien-			
tation in Pharmacy, The (E)	200	Weiss, Benjamin—	
New Drugs of 1959	46	Synthetic Analgesics, Part I,	
Our Most Critical Need (E)	120	Diphenylpropylamines (B) .	307

SUBJECT INDEX, 1960

(A)—Abstract

(B)—Book Review

(E)—Editorial

	PAGE		PAGE
Acetazolamide	17	Anturan	53
Acetylcholine	129	Aristocort Syrup	69
Actasal Pediatric Drops	52	Arlidin	59
Actase	72	Arteriosclerosis	395
ACTH/Cortisone	147	Arthropan Liquid	52
A Decade of Chemotherapeutic Management of Chronic Granu- locytic and Chronic Lympho- cytic Leukemia	145	Ascorbic Acid	363, 441
Ademol	54	Aspartocin	48
Airborne Bacteria Transmission in Hospital Wards (A)	421	Aspirin (E)	84
Aldactone	55	Atherosclerosis	390
Aldosterone	55	Aureomycin	47
Alpha Chymar	73	BAL	334
Alphaprodine	226	B. C. M.	234
Altafur	49	Benzimidazole Mustard	235
A. M. A. and Proprietary Drugs (E)	353	Benzquinonium Chloride	129
American Society of Pharma- cognosy	271	Benzthiazide	55
Amine Oxidase Inhibitors	96	Bevitam	71
Amino Carboxylic Acids	328	Biochemical Preparations, Vol- ume 7 (B)	386
Amotriphene	59	Bioflavonoids	363
Amphetamine	61, 96, 360	Bioflavonoid Therapy (A)	108
Amphotericin B (A)	30	Biophysical Science: A Study Program (B)	273
Analgesic Substances	202	Bransky, Joseph M., Retirement	352
Anectine	129	Calcium Acetylsalicylate Carba- mide	52
An Example for Others (E) ...	250	Calurin	52
An Introduction to Pharmacol- ogy (B)	306	Carbopol 934 as Ephedrine Sul- fate Jelly Base	383
Anorexiants	96	Cardiografin	75
A Note on the Solubility of Salicylic Acid in Isopropanol- Water Solutions at 25°C. ...	406	Carisoprodol	64, 141
Antibacterials (A)	34	Caytine	61
Antibiotic Activity (A)	109	Chemipen	47
Antibiotics	147	Chemotherapy	338
Antidepressives	96	Chlorambucil	231
		Chlordantoin	50
		Chlorphenoxamine Hydrochloride	62
		Chlortetracycline	47
		Chlorothiazide	18, 53
		Chlorpromazine	359

	PAGE		PAGE
Cholesterol	402	Diethylpropion	61
Choline Salicylate	52	Dimethisterone (A)	109
Cholinesterase Inhibitors	125	Disamide (A)	110
Cholografin	75	Diuretic Agents	5
Chronology of Atherosclerosis ..	390	Drug Counterfeiting (E)	313
Chymar Buccal	73	Drug or Beverage? (E)	94
Coal Tar Ointments (A)	31	Drug Research Today (E)	44
Colchicines	147	Drugs, Their Nature, Action and	
Colistin	49	Use (B)	78
Coly-mycin M	49	Duografin	75
Coly-mycin S	49	Durabolin	70
Combistix Reagent Strips	76	Edathamil Disodium	76
Composition Studies on Tobacco	165	Edema (A)	421
Connective Tissue, Thrombosis		Edrophonium	136, 143
(B)	39	EDTA	334
Connective Tissue, Thrombosis		Elastase	73
and Atherosclerosis (B)	39	Embichin	234
Constipation Control in Subma-		Emulsions	315
riners (A)	422	Endoxan	51
Contribution to Western Penn-		Endrate Disodium	76
sylvania Pharmacy, A (B) ...	37	Enovid	70
CR-Test	76	Ephedrine Sulfate Jelly	383
Curare	129	Epinephrine	357
Cyanocobalamin	448	Epoxypropidine (A)	113
Cyclophosphamide	50	Erythromycin Salts and Tri-	
Cytoxan	50	cetyloleandomycin (A)	112
Deanol	360	Eserine	129
Decade of Chemotherapeutic		Evaluation of Analgesic Sub-	
Management of Hodgkin's Dis-		stances	202
ease	27	Evaluation of Diuretic Agents .	5
Decade of Management of Poly-		Evaluation of Drugs Affecting	
cythemia Through Chemo-		Neuro-Muscular Transmission	122
therapy	338	Expansion of Drug Certification	
Decamarium Bromide	56	by the FDA (E)	431
Decamethonium	129	Exponential De-Emulsification	
Declomycin	47	and Stability of Pharmaceu-	
De-Emulsification	315	tical Emulsions	315
Deepain	205	Fat Consumption and Coronary	
Degranol	234	Disease: The Evolutionary An-	
Depinar	71	swer to the Problem (B) ...	276
Depo-Medrol	69	Feosol Spansules	72
Desoxyephedrine	61	Fervenuin	48
Dexedrine	224	Fibrinolysin	72
Dextromoramide	52	Fibrinuclease	73
Diamino-Diphenyl Sulfone (A)	239	Figures That Count: Mathe-	
Diamox	56	matics for Nurses (B)	275

	PAGE		PAGE
Flaxedil	129	JB-591	58
Floropryl	129	Konakion	71
Flumethiazide	54	Lavandula Multifida L.	252
Fluorometholone	69	Leukemia	145, 416
Fluphenazine	63	Lipide Chemistry (B)	423
Folic Acid Antagonists ..147, 341,	415	LSD	360
Fortespan	71	Lymphocytic Leukemia	145
Fulvicin	48	Lymphosarcoma	296, 417
Furaltadone	49	Malignant Blood Diseases	415
GABA	358	Marplan	67
Gallamine Triethiodide	129	Maxipen	47
Gargon	49	Mecostin	129
Gastrografin	75	Medieval and Renaissance Medi-	
Generic Name Pitfall (E)	2	cine (B)	38
Genus Lavandula	252	Medrol	59
GG-Test	76	Medroxyprogesterone Acetate ..	70
Gibberellic Acid	98	Mellaril	63
Government Purchases of Drugs		MER-29	60
Overseas (E)	280	6-MP	147
Granulocytic Leukemia	145	Meralluride	17
Griffith, Ivor—Remington Med-		Mestinon	129
alist 1960	428	Mephesisin	141
Grifulvin	48	Methazolamide	56
Griseofulvin	48	Methoden der Organischen	
Guanethidine	57, 385	Chemie (Houben-Weyl), Vol.	
Herbs	178	1, Part 2 (B)	195
Heterocyclic Chemistry (B) ...	456	Methoden der Organischen	
Hexadimethrine Bromide	76	Chemie (Houben-Weyl), Vol.	
Histamine	357	5, Part 4 (B)	305
Histological Studies of the Genus		Methohexital	76
Lavandula	252	A-Methopterin	51
Hodgkin's Disease ...27, 231, 295,	417	Methotrexate	51
Humatin	49	Methoxypromazine Maleate	63
Humorsol	56	Methylphenidate	360
Hydrochlorothiazide	19, 421	Metubine	129
Hydroxy Carboxylic Acids	325	Modern Pharmacology and	
Iloson	48	Therapeutics (B)	196
Imipramine	360	Mornidine	65
Imipramine Hydrochloride	68	Multiple Antigen (A)	34
Immunization (A)	34	Multiple Myeloma	298, 418
Interneuronal Blocking Agents .	95	N-Mustards	147
Ismelin	57	Mycosis Fungoides	298, 418
Isocarboxazid	67	Myleran	147
Isordil	58	Myordil	59
Isosorbide Dinitrate	58	Mytolon	129
Isosuprine Hydrochloride	58	Nandrolone Phenpropionate	70

	PAGE		PAGE
Naphazoline Hydrochloride	60	Pantothenic Acid	446
Nardil	66	Pentothal	141
National Association of Boards of Pharmacy	409	Permitil	63
Naturetin	54	Pharmacist and Research	158
Need for Mutual Support (E) ..	156	Pharmacology in Nursing (B) ..	386
Need for Professional Orienta- tion in Pharmacy (E)	200	Phenazocine Bromide	51
Neostigmine	129	Phenelzine	66
Neptazane	56	Phenistix	75
Neurochemistry of Nucleotides and Amino Acids (B)	307	Phenmetrazine	61
Neurohormone-Phrenotropic Drug Activity Correlates	355	Phenothiazine Derivatives	94
Neurosedatives	95	Phenoxene	62
New and Non-Official Drugs, 1959 (B)	274	Phenylbutazone	52, 225
New Drugs of 1959	46	Phospholine Iodide	56
New Nitrogen Mustards in the Management of Hodgkin's Dis- ease	231	Phosphoramides in the Manage- ment of Lymphomas and Reticuloses	295
Nialamide	67	Phrenotropic Drugs	360
Niamid	67	Physostigmine	128
NIH-7519	51	Pipamazine	65
Nitrogen Mustards	231, 340	Piperidine Derivatives	96
Non-Barbiturate Sedative-Hyp- notics	95	Piperilate HCL	64
Norethindrone	70	Pipradrol	360
Norethynodrel	70	Plant Science Seminar	271
Norlutin	70	Plurionics as a Suppository Base ..	301
Novoembichin	234	Polybrene	76
Nutrition and Atherosclerosis (B)	38	Polycythemia	338, 418
Nylidrine Hydrochloride	59	Polycythemia Vera	298
Objectives of the N. A. B. P. ..	409	Polypeptides	283
Organic Reactions, Volume X (B)	423	Polyphosphates	326
Organic Syntheses, Vol. 39 (B) ..	37	Powers, Justin L., cover dedica- tion	312
Ostensin	57	Prinadol	51
Otrivin	60	Privine	60
Our Most Critical Need (E) ...	120	Progesterin	70
Oxidation—Reduction Potentials of Organic Systems (B)	241	Proteins	283
Oxylone	69	Project HOPE (E)	388
P-32	147	Prolixin	63
Palfum	52	Prostigmin	129
		Provera	70
		Psychotherapeutic Drugs	86
		Quadrigen	74
		Radiophosphorus	338
		Rauwolfia Preparations	95
		Rela	64
		Remington Medalist for 1960 and His Co-Workers	428

	PAGE		PAGE
Reserpine	359	TEM	147
Resprogen Injection	74	Tepanil	61
Reticuloendotheliosis	418	Terramycin I. M.	47
Reticulum-Cell Sarcoma	297	Tetravax	74
Retrogratin	75	Thiamine	437
Riboflavin	440	Thioridazine Hydrochloride ..	63
Rifomycin B	48	Thiostrepton	49
Salicylic Acid Solubility	406	Tigan	66
Saluron	54	Tobacco	165
S. A. Vite	72	Tofranil	68
Sequence of Amino-Acids in Proteins and Polypeptides ..	282	Tolbutamide Therapy (A)	33
Sequestration and Its Applica- tions	325	Tranquilizers	63
Shakers as Pioneers in the Amer- ican Herb and Drug Industry	178	Triamcinolone	69
Sinografin	75	Triburon	50
Smallpox Vaccine Avianized ...	74	Triclobonium Chloride	50
SKF-385A	68	Triethylene Melamine	339
Sodium Diatrizoate	75	Trilafon	63
Solu-Medrol	69	Trimethidinium Methosulfate ..	57
Soma	64, 141	Trimethoxybenzamide Hydro- chloride	66
Spironolactone	55, 421	Triparanol	59
Sporostacin Cream	50	Tylosin	49
Stability Problems with Some Vitamins in Pharmaceuticals ..	433	Unique Place of Folic Acid An- tagonists in Controlling Malign- ant Blood Diseases	415
Steroid Activity (A)	239	Unpublished Abstracts of Ar- ticles on Pharmaceutical Sub- jects, Series 8 (B)	115
Story of Chemistry (B)	347	Uracil Mustard	235
Streptozotocin	48	Urdang, George, In Memoriam (E)	248
Subsidia Pharmaceutica 1959 (B)	36	Urethan	147
Subsidia Pharmaceutica and Index Nominum 1960 (B) ..	304	Vanillic Diethylamide	77
Succinylcholine	129	Varidase Buccal	73
Sucostrin	129	Vasodilan	58
Sulfinpyrazone	53	Velacycline	47
Supain	204	Vitamin A	115, 434
Suppository Base	301	Vitamin B ₁₂	71, 444
Sycotrol	64	Vitamin K (A)	194
Syncillin	46	Vitamin K ₁	71
Syncurine	129	Vitamin-Mineral Compositions ..	433
Syntetrin	47	Vitamin P	363
Synthetic Analgesics, Part I, Diphenylpropylamines (B) ..	307	Vitamins	433
Tensilon	136	Water Absorption by Petrolatum Ointment Bases (A)	111
Tentone	63	Xylometazoline Hydrochloride ..	60
Tenuate	61	Zolyse	73

THE AMERICAN
JOURNAL OF PHARMACY
and The Sciences Supporting Public Health

A RECORD OF THE PROGRESS OF PHARMACY AND THE
ALLIED SCIENCES

Since 1825

Published Monthly by

THE
PHILADELPHIA COLLEGE OF
PHARMACY AND SCIENCE
43d Street Kingsessing and Woodland Avenues
Philadelphia 4, Pennsylvania

LINWOOD F. TICE, Ph. G., M. Sc., D. Sc., Editor
Kenneth Avis, M. Sc., D. Sc., Editorial Assistant
Margaret Mary Culin, Editorial Assistant
John E. Kramer, B. Sc., Business Manager

COMMITTEE ON PUBLICATION

Louis Gershenfeld, P. D., Ph. M., D. Sc., Chairman
Mitchell Bernstein, P. D., M. D., F. A. C. P.
E. Fullerton Cook, P. D., Ph. M., D. Sc.
Marin S. Dunn, A. M., Ph. D.
Joseph W. E. Harrison, P. D., Sc. D.
Ivor Griffith, P. D., Ph. M., D. Sc., F. R. S. A., ex officio

VOLUME 132

1960

PHILADELPHIA, PA.

Printed at
Philadelphia, Pa.

NEW!

PLANOLAR

TRADEMARK

TABLETS

FOR 2-PRONGED ATTACK ON RHEUMATOID ARTHRITIS



*Immediate
Relief of Pain*

*Cumulative
Antirheumatic
Anti-inflammatory
Treatment*

With new Planolar, patients with rheumatoid arthritis can obtain immediate relief of pain and discomfort and, at the same time, begin specific long-term therapy. This new tablet combines the effective antirheumatic and anti-inflammatory agent, Plaquenil® (60 mg.), with the time-tested analgesic, aspirin (300 mg.). The usual adult dose is 2 tablets two or three times daily.

Years of use have demonstrated that Plaquenil is highly successful in the treatment of patients with rheumatoid arthritic conditions. The improvement obtained with

Plaquenil is generally well tolerated and its use can be continued almost indefinitely in most patients. Aspirin, the other ingredient in Planolar, continues to be the analgesic of choice in the control of pain in rheumatoid disease.

The vast rheumatoid arthritis market indicates heavy prescription ahead for Planolar. Order full stocks now. Planolar will be heavily promoted to the profession through journal advertising, direct mail and detailing to physicians in your area.

PLANOLAR IS AVAILABLE IN BOTTLES OF 100 TABLETS.

Winthrop

LABORATORIES
New York 18, N.Y.



high
potency
vitamin-
mineral
formula
for
active
people

**MYADEC
CAPSULES**

When bodily stresses increase vitamin and mineral requirements, you can recommend MYADEC with complete confidence. MYADEC helps prevent vitamin-mineral deficiencies by providing comprehensive nutritional supplementation—one capsule daily supplies therapeutic doses of nine important vitamins plus significant quantities of eleven minerals and trace elements. Be sure you have an adequate stock of MYADEC for display. Supplied: Bottles of 30, 100, and 200.

PARKE-DAVIS & COMPANY
Grand St. Bridge

PARKE-DAVIS

American Journal of Pharmacy

The American Journal of Pharmacy is the oldest continuously published scientific periodical of its kind in America, having been established by the Philadelphia College of Pharmacy in 1825. After the original issue there were three other preliminary numbers until 1829, when regular publication began. From then until 1852 four issues were published annually, with the single exception of 1847, when an additional number appeared. Six issues a year were printed from 1853 to 1870, at which time the Journal became a monthly publication.

Former Editors of the Journal have been: Daniel B. Smith, 1825-1828; Benjamin Ellis, 1829-1831; Robert E. Griffith, 1831-1836; Joseph Carson, 1836-1850; William Procter, Jr., 1850-1871; John M. Maisch, 1871-1893; Henry Trimble, 1893-1898; Henry Kraemer, 1898-1917; George M. Beringer, 1917-1921, and Ivor Griffith, 1921-1941.

Established and maintained as a record of the progress of pharmacy and the allied sciences, the Journal's contents and policies are governed by an Editor and a Committee on Publications elected by the members of the College.

Manuscripts should be sent to the Editor, who does not assume any responsibility in connection with the views or investigations of contributors of accepted manuscripts, other than to exercise general care in selection.

Contributors are allowed a reasonable number of copies of this Journal, free of charge, if applied for when the proof is returned.

Reprints, if desired, should be ordered when the proof is returned. The table below shows the *approximate* cost of reprints, the make-up of the pages to be identically the same as in the Journal. The actual cost may vary from the figures given, and will depend upon the amount of presswork, paper, binding, and other factors. Reprints containing half-tones may be expected to cost somewhat more than the rates given.

	2 pp.	4 pp.	8 pp.	16 pp.	COVERS WITH TITLES
50 copies.....	\$ 4.50	\$10.00	\$16.25	\$27.50	50 copies..... \$ 7.50
100 "	7.50	13.75	21.25	40.00	100 " 12.50
250 "	10.00	17.50	27.50	53.75	250 " 17.50
500 "	15.00	25.00	35.00	68.75	500 " 26.25



C A N CAN BE CURED

More than one million Americans are living proof. Remember . . . your contributions helped save many of these lives. Your continuing contributions are needed to help discover new cures and, ultimately, the prevention of cancer itself • Remember, too, if you delay seeing your physician, you drastically cut your chances of cure. Annual checkups are the best way of detecting cancer in time • Guard your family! Fight cancer with a checkup and a check.

AMERICAN CANCER SOCIETY

